

CHECK-UP CARE

You would do anything to keep your child happy, healthy and protected against diseases. We would, too! To optimize your child's growth and development — and ensure your child is fully immunized — please schedule well child visits at each of the following age milestones:

| | Measurements | Screenings | Care | Tests & Labs | Immunizations |
|-------------|--------------------------------------|--|-------------------------|-------------------------|---|
| Newborn | Length Weight Headsizes | | | | |
| 1 month | Length Weight Headsizes | Maternal Depression | | | |
| 2 months | Length Weight Headsizes | Maternal Depression Child Development | | | DTaP-IPV-HepB ¹ , Hib, PCV13, RV1 |
| 4 months | Length Weight Headsizes | Maternal Depression Child Development | | | DTaP-IPV-HepB ¹ , Hib, PCV13, RV1 |
| 6 months | Length Weight Headsizes | Maternal Depression Child Development | Fluoride Dental Varnish | | DTaP-IPV-HepB ¹ , PCV13, Influenza ² |
| 9 months | Length Weight Headsizes | Child Development | Fluoride Dental Varnish | | Influenza ² |
| 12 months | Length Weight Headsizes | Child Development | Fluoride Dental Varnish | Hemoglobin Lead Test | MMR, VAR, HepA, Influenza ² |
| 15 months | Length Weight Headsizes | Child Development | Fluoride Dental Varnish | | DTaP, Hib, PCV13, Influenza ² |
| 18 months | Length Weight Headsizes | Child Development | Fluoride Dental Varnish | | HepA, Influenza ² |
| 2 years | Height Weight Headsizes BMI | Child Development | Fluoride Dental Varnish | Lead Test | Influenza ² |
| 2-1/2 years | Height Weight BMI | Child Development | Fluoride Dental Varnish | | Influenza ² |

| | Measurements | Screenings | Care | Tests & Labs | Immunizations |
|-----------------|--|---|-------------------------|---|---|
| 3 years | Height / Weight BMI Blood pressure | Child Development | Fluoride Dental Varnish | Vision | Influenza ² |
| 4 years | Height / Weight BMI Blood pressure | Child Development | Fluoride Dental Varnish | Vision Hearing | Influenza ² , DTaP-IPV, MMR-VAR (these combination vaccines can be given and age 4 or 5) ¹ |
| 5 years | Height / Weight BMI Blood pressure | Child Development | Fluoride Dental Varnish | Vision Hearing | Influenza ² , DTaP-IPV, MMR-VAR (these combination vaccines can be given and age 4 or 5) ¹ |
| 6-10 years* | Height / Weight BMI Blood pressure | Social & Emotional Development | | Vision Hearing Lipid Profile ³ | Influenza ² |
| 11 years | Height / Weight BMI Blood pressure | Social & Emotional Development | | Vision Hearing | Tdap, MCV4, HPV ⁴ , Influenza ² |
| 12-15 years* | Height / Weight BMI Blood pressure | Social & Emotional Development Depression | | Vision Hearing | Influenza ² |
| 16 years | Height / Weight BMI Blood pressure | Social & Emotional Development Depression | | Vision Hearing | MenB ⁵ , MCV4, Influenza ² |
| 17 years | Height / Weight BMI Blood pressure | Social & Emotional Development Depression | | Vision Hearing | Influenza ² |
| 18 years | Height / Weight BMI Blood pressure | Depression | | Vision Hearing Lipid Profile HIV | Influenza ² , Tdap |
| 19 years+ | Height / Weight BMI Blood pressure | Depression | | Hearing Vision | Influenza ² |

| Notes | |
|--|--|
| ¹ Combination Vaccines - Pediarix (DTaP-IPV/HepB), Kinrix (DTaP-IPV), ProQuad (MMRV=MMR and VAR) | ⁴ HPV (Gardasil) - Routinely started at age 11 years but can be started at age 9 years. If started before age 15 years, it is a 2 dose series given 6 months apart. |
| ² Influenza Vaccine - Given once each year because influenza strains can change from year to year | ⁵ MenB (Bexsero) - Given as a 2 dose series at least 1 month apart |
| ³ Done one time | *Schedule visits annually |